

Cheap Nursing CEUs

COVID-19: An Update for 2021

1. A patient has questions regarding COVID-19. Which of the following educational points should be expressed to the patient?

- A. COVID-19 can spread through person-to-person contact.
 - B. COVID-19 is primarily transmitted through the air.
 - C. COVID-19 is only transmitted by contact with infected blood.
 - D. COVID-19 is only transmitted via contaminated food.
-

2. According to the course, how long can the COVID-19 virus live on surfaces?

- A. Up to 7 days
 - B. Up to 10 days
 - C. Up to 14 days
 - D. Up to 28 days
-

3. According to the course, what is the suggested incubation period for COVID-19?

- A. 1 - 14 days
 - B. 14 - 18 days
 - C. 18 - 20 days
 - D. 20 - 30 days
-

4. According to the course, which of the following may be a symptom of COVID-19?

- A. Seizures
 - B. Weight gain
 - C. Fever
 - D. Rash
-

5. A health care professional has questions regarding antigen tests for COVID-19. Which of the following informational points of interest should be communicated to the health care professional?

- A. Antigen tests should not be used on individuals over the age of 65.
 - B. Antigen tests should not be used on individuals under the age of 65.
 - C. Antigen tests may not be as sensitive as NAATs, and the results of an antigen test may need to be confirmed with a NAAT.
 - D. Antigen tests are more sensitive than NAATs.
-

6. According to the course, what is the recommended dose for the Pfizer-BioNTech COVID-19 vaccine?

- A. 0.2 mL**
 - B. 0.3 mL**
 - C. 0.4 mL**
 - D. 0.5 mL**
-

7. A 48-year-old female patient has questions regarding the Moderna COVID-19 vaccine. Which of the following educational points should be expressed to the patient?

- A. The vaccine should be administered in 2-dose series separated by 18 days.**
 - B. The vaccine should be administered in 2-dose series separated by 28 days.**
 - C. The vaccine cannot be offered to individuals that have a history of prior symptomatic or asymptomatic COVID-19 virus infection.**
 - D. The vaccine cannot be offered to individuals, over the age of 40, that have a history of prior symptomatic or asymptomatic COVID-19 virus infection.**
-

8. A health care professional has questions regarding bamlanivimab. Which of the following informational points of interest should be communicated to the health care professional?

- A. Bamlanivimab should be considered the standard of care for the treatment of patients with COVID-19.**
 - B. Bamlanivimab should be considered the standard of care for the treatment of patients over the age of 18 with COVID-19.**
 - C. Bamlanivimab must be administered by IV infusion.**
 - D. Bamlanivimab must be administered by IM injection.**
-

9. What is the recommended adult Day 1 dose of remdesivir for the treatment of COVID-19?

- A. 100 mg**
 - B. 200 mg**
 - C. 400 mg**
 - D. 500 mg**
-

10. A 24-year-old male patient has questions regarding variant B.1.1.7. Which of the following educational points should be expressed to the patient?

- A. Variant B.1.1.7 does not affect individuals under the age of 25.**
- B. Variant B.1.1.7 emerged independently in South Africa.**

C. Variant B.1.1.7 spreads easily and quickly when compared to other COVID-19 variants.

D. Variant B.1.1.7 has not been detected in the U.S.

11. A fellow health care professional has questions regarding hand hygiene. Which of the following informational points of interest should be communicated to the health care professional regarding hand hygiene?

A. A health care professional should engage in hand hygiene after touching a patient.

B. The duration of the entire handwashing procedure with soap and water should last between 10 - 20 seconds.

C. The duration of the entire handwashing procedure with soap and water should last between 5 - 10 seconds.

D. At the end of a hand hygiene procedure with an alcohol-based formulation, health care professionals should dry their hands with a towel.

12. A health care professional has questions regarding hand hygiene with an alcohol-based formulation. Which of the following informational points of interest should be communicated to the health care professional?

A. The duration of the entire procedure should last between 5 - 10 seconds.

B. The duration of the entire procedure should last between 10 - 15 seconds.

C. The duration of the entire procedure should last between 15 - 18 seconds.

D. The duration of the entire procedure should last between 20 - 30 seconds.

13. Which of the following statements is most accurate?

A. All individuals with signs and symptoms of a respiratory infection (excluding health care professionals) should be instructed to cover the mouth and nose with their hands when coughing or sneezing.

B. All individuals with signs and symptoms of a respiratory infection (including health care professionals) should be instructed to cover the mouth and nose with their hands when coughing or sneezing.

C. All individuals over the age of 65 with signs and symptoms of a respiratory infection should be instructed to cover the mouth and nose with their hands when coughing or sneezing.

D. All individuals with signs and symptoms of a respiratory infection (including health care professionals) should be instructed to cover the mouth and nose with a tissue when coughing or sneezing.

14. A health care professional has questions regarding safe injection practices. Which of the following informational points of interest should be communicated to the health care professional?

A. Health care professionals should always unwrap syringes prior to the time of use for

all patients.

B. Health care professionals should always unwrap syringes prior to the time of use for patients over the age of 65.

C. Health care professionals may administer medications from the same syringe to multiple patients.

D. Health care professionals should never administer medications from the same syringe to multiple patients.

15. A fellow health care professional has questions regarding contact precautions. Which of the following informational points of interest should be communicated to the health care professional regarding contact precautions?

A. If contact precautions may be required, patients with symptoms of an active infection should be placed in a private room until it is determined if contact precaution are necessary.

B. Signs indicating patients are on contact precautions should not be posted outside patient rooms.

C. PPE is not necessary if contact precautions are required for a patient.

D. Patient transport and movement should not be limited.

16. Which of the following statements regarding airborne precautions is most accurate?

A. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should place patients in a private room with the door open.

B. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should only place patients over the age of 65 in a private room with the door open.

C. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should place patients in a private room with the door closed.

D. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should only place patients over the age of 65 in a private room with the door closed.

17. A 52-year-old female patient requires droplet precautions. How can health care professionals ensure droplet precautions are applied to the patient?

A. Ensure the patient wears a face shield and goggles at all times.

B. Ensure the patient wears a facemask, face shield, and goggles at all times.

C. Allow the patient to exit the room for periods of 20 - 90 minutes.

D. Limit the transport and movement of patients outside of their rooms to medically-necessary purposes.

18. A health care professional has questions regarding the Interim Infection Prevention and Control Recommendations. Which of the following informational points of interest should be communicated to the health care professional?

- A. Facemasks are preferred over cloth face masks for health care professionals.**
 - B. Cloth masks are preferred over facemasks for health care professionals.**
 - C. Facemasks are not considered to be PPE.**
 - D. Cloth masks are considered to be PPE.**
-

19. An adult individual enters a health care facility. The adult individual is carrying a child under the age of two. A health care professional observes that the individual and the child are not wearing masks. According to the Interim Infection Prevention and Control Recommendations, how should the health care professional proceed?

- A. Place a facemask on the adult individual and the child.**
 - B. Place a respirator on the adult individual and a facemask on the child.**
 - C. Offer the adult individual and the child a facemask or mask.**
 - D. Offer the adult individual a facemask or mask, and note that facemasks and cloth masks should not be placed on young children under age two.**
-

20. A health care professional has questions regarding COVID-19 screening. Which of the following informational points of interest should be communicated to the health care professional?

- A. Screening for fever and symptoms of COVID-19 should be incorporated into the daily assessments of all admitted patients.**
 - B. Screening for fever and symptoms of COVID-19 should only be incorporated into the daily assessments of patients over the age of 65.**
 - C. Screening for fever and symptoms of COVID-19 should only be incorporated into the daily assessments of patients under the age of 18.**
 - D. Screening for fever and symptoms of COVID-19 should not be incorporated into the daily assessments of patients.**
-