

# Cheap Nursing CEUs

## Fragile X Syndrome

1. Fragile X syndrome is more common in females.

- A. True
  - B. False
- 

2. Which of the following may be considered a physical symptom associated with Fragile X syndrome?

- A. Large head
  - B. Large teeth
  - C. Small ears
  - D. Wide face
- 

3. Which of the following may be considered a behavioral, social, and emotional symptom associated with Fragile X syndrome?

- A. Hyperactivity
  - B. An inability to learn new skills
  - C. An inability to efficiently or effectively solve problems
  - D. Stuttering
- 

4. Which of the following statements is most accurate?

- A. Often individuals show symptoms of Fragile X syndrome by 2 months to 4 months of age.
  - B. Often individuals show symptoms of Fragile X syndrome by 6 months to 10 months of age.
  - C. The physical symptoms associated with Fragile X syndrome typically develop when an individual is about 6 months of age.
  - D. The physical symptoms associated with Fragile X syndrome typically develop when an individual goes through puberty.
- 

5. Which of the following may be considered an early sign/symptom of Fragile X syndrome?

- A. Begins to speak early
- B. Learns to walk early
- C. Displays unusual reactions to the way things sound, smell, taste, look, or feel

**D. Enlarged testicles in males**

---

**6. In order to be diagnosed with an intellectual disability, an individual must have both a significantly low IQ and considerable problems in everyday functioning.**

- A. True**
  - B. False**
- 

**7. A patient has questions regarding Celexa. Which of the following educational points should be expressed to the patient?**

- A. Celexa is a SSRI.**
  - B. Celexa is an antipsychotic agent.**
  - C. Celexa is not associated with nausea or dry mouth.**
  - D. A typical dose of Celexa is 100 mg daily.**
- 

**8. A patient has questions regarding Concerta. Which of the following educational points should be expressed to the patient?**

- A. Concerta is a SSRI.**
  - B. Concerta is an antipsychotic agent.**
  - C. A typical starting dose of Concerta for children and adolescents is 80 mg once daily.**
  - D. A typical starting dose of Concerta for children and adolescents is 18 mg once daily.**
- 

**9. What is a typical dose of Adderall XR for individuals ages 6 - 17?**

- A. 10 mg once daily**
  - B. 50 mg once daily**
  - C. 100 mg once daily**
  - D. 100 mg twice daily**
- 

**10. A genetic counselor is an individual who specializes in genetics and provides patients with counseling, education, and assistance in making decisions regarding genetic-related health care.**

- A. True**
  - B. False**
-