Cheap Nursing CEUs

Fragile X Syndrome

1. Fragile X syndrome is more common in females.

A. True
B. False

---

2. Which of the following may be considered a physical symptom associated with Fragile X syndrome?

A. Large head
B. Large teeth
C. Small ears
D. Wide face

---

3. Which of the following may be considered a behavioral, social, and emotional symptom associated with Fragile X syndrome?

A. Hyperactivity
B. An inability to learn new skills
C. An inability to efficiently or effectively solve problems
D. Stuttering

---

4. Which of the following statements is most accurate?

A. Often individuals show symptoms of Fragile X syndrome by 2 months to 4 months of age.
B. Often individuals show symptoms of Fragile X syndrome by 6 months to 10 months of age.
C. The physical symptoms associated with Fragile X syndrome typically develop when an individual is about 6 months of age.
D. The physical symptoms associated with Fragile X syndrome typically develop when an individual goes through puberty.

---

5. Which of the following may be considered an early sign/symptom of Fragile X syndrome?

A. Begins to speak early
B. Learns to walk early
C. Displays unusual reactions to the way things sound, smell, taste, look, or feel
D. Enlarged testicles in males

---

6. In order to be diagnosed with an intellectual disability, an individual must have both a significantly low IQ and considerable problems in everyday functioning.

A. True
B. False

---

7. A patient has questions regarding Celexa. Which of the following educational points should be expressed to the patient?

A. Celexa is a SSRI.
B. Celexa is an antipsychotic agent.
C. Celexa is not associated with nausea or dry mouth.
D. A typical dose of Celexa is 100 mg daily.

---

8. A patient has questions regarding Concerta. Which of the following educational points should be expressed to the patient?

A. Concerta is a SSRI.
B. Concerta is an antipsychotic agent.
C. A typical starting dose of Concerta for children and adolescents is 80 mg once daily.
D. A typical starting dose of Concerta for children and adolescents is 18 mg once daily.

---

9. What is a typical dose of Adderall XR for individuals ages 6 - 17?

A. 10 mg once daily
B. 50 mg once daily
C. 100 mg once daily
D. 100 mg twice daily

---

10. A genetic counselor is an individual who specializes in genetics and provides patients with counseling, education, and assistance in making decisions regarding genetic-related health care.

A. True
B. False

Copyright © 2026 Cheap Nursing CEUs

Visit us at https://www.cheapnursingceus.com