

# Cheap Nursing CEUs

## HIV Treatment

1. Which of the following statements regarding antiretroviral therapy (ART) is most accurate?

- A. ART is recommended for HIV-infected individuals for the prevention of transmission of HIV.
  - B. Studies have shown ART cannot prevent the transmission of HIV.
  - C. Studies have shown ART cannot improve HIV patients' immune system function.
  - D. None of the above
- 

2. A 48 year-old male, HIV patient presents to your hospital with an acute infection. He is currently on no medications and his CD4 count is below 350 cells/ $\mu$ L. Which of the following recommendations can be made to reduce HIV-related morbidity?

- A. Initiate an ART treatment regimen when the patient's CD4 count is below 300
  - B. Initiate an ART treatment regimen when the patient's CD4 count is below 200
  - C. Initiate an ART treatment regimen as soon as possible
  - D. None of the above
- 

3. You are taking an HIV patient's history upon admission. It is important to gather information on which of the following?

- A. Medication allergies
  - B. HIV RNA (viral load) history
  - C. Current medications, including herbal preparations
  - D. All of the above
- 

4. Which of the following should be included in the first step to initiating a patient on an ART regimen?

- A. Select the appropriate medications
  - B. Review the medication side effects with the patient
  - C. Have a detailed discussion with the patient about his or her readiness to commit to a medication regimen
  - D. None of the above
- 

5. ART medications cannot be initiated after the presence of an opportunistic infection.

- A. True

**B. False**

---

**6. Which of the following medications are considered to be a NRTI fixed-dose, combination formulation?**

- A. Truvada**
  - B. Epzicom**
  - C. Kaletra**
  - D. Both A and B**
- 

**7. Preferred INSTI based regimens may include Raltegravir.**

- A. True**
  - B. False**
- 

**8. Your patient is initiated on ART medications. Which of the following should be monitored?**

- A. Adverse reactions**
  - B. CD4 count**
  - C. Blood pressure**
  - D. All of the above**
- 

**9. "Antiretroviral (ARV) medications can be given in two-drug regimens to decrease the potential for developing of resistance."**

- A. True**
  - B. False**
- 

**10. A 32 year-old women is admitted to your unit. She is currently experiencing ART treatment failure. Which of the following factors could have lead to her treatment failure?**

- A. Poor ARV medication adherence**
  - B. ARV resistance**
  - C. CD4 count elevation**
  - D. Both A and B**
- 

**11. Combination ARV regimens are recommended for all HIV-infected women during pregnancy, regardless of CD4 count.**

- A. True**
  - B. False**
-

**12. Which of the following ARV medications can be used during pregnancy?**

- A. Tenofovir**
  - B. Kaletra**
  - C. Nevirapine**
  - D. All of the above**
- 

**13. "HIV-infected pregnant women who have not received ARV prior to labor should be given ZDV as a continuous infusion during labor."**

- A. True**
  - B. False**
- 

**14. What is the recommended infant ZDV dose?**

- A. 4 mg/kg/dose PO QID**
  - B. 4 mg/kg/dose PO BID**
  - C. 8 mg/kg/dose PO BID**
  - D. 8 mg/kg/dose PO TID**
- 

**15. Which of the following should be evaluated during every medical visit by pregnant women with HIV?**

- A. Blood pressure**
  - B. Intimate-partner violence**
  - C. HIV antibody test**
  - D. Both A and B**
- 

**16. Which of the following immunizations are recommended for pregnant women with HIV infection?**

- A. Varicella**
  - B. Measles/Mumps/Rubella**
  - C. Tetanus-diphtheria-pertussis**
  - D. Both A and B**
- 

**17. "Heterosexual transmission of HIV is more efficient from man to woman than from woman to man."**

- A. True**
  - B. False**
-

18. Which of the following ARV adverse effects may be more severe in women?

- A. Anemia
  - B. Bone loss
  - C. Lactic acidosis
  - D. All of the above
- 

19. A 36 year-old male patient is admitted to your unit for palliative HIV care. Which of the following treatment strategies may be used as supportive care?

- A. Pain management
  - B. Nausea management
  - C. Depression treatment
  - D. All of the above
- 

20. Megestrol acetate may improve appetite and fatigue in HIV patients.

- A. True
  - B. False
- 

21. Which of the following medications may be used to treat HIV related constipation?

- A. Senna
  - B. Docusate
  - C. Zofran
  - D. Both A and B
- 

22. You are asked to educate your HIV patient on advance care planning, what should you include?

- A. Advance directive information
  - B. Health care proxy information
  - C. Pain management information
  - D. All of the above
- 

23. You are asked to assess your patient's ART medication adherence, what questions should you ask your patient?

- A. What HIV medications do you take and what are their dosages?
  - B. How many doses of your HIV medication have you missed in the past 72 hours?
  - C. Are you comfortable taking medications in front of others?
  - D. All of the above
-

**24. Advanced HIV infection does not impact patient medication adherence.**

- A. True**
  - B. False**
- 

**25. Which of the following strategies can improve ART medication adherence?**

- A. Anticipate and treat adverse effects**
  - B. Simplify medication regimens**
  - C. Provide substance abuse recourses**
  - D. All of the above**
- 

**Copyright © 2023 Cheap Nursing CEUs**

**Visit us at <https://www.cheapnursingceus.com>**