

Cheap Nursing CEUs

Obesity

1. What is the most common diagnostic tool currently used for identifying obesity according to the course content, and why is there a shift towards considering other measures?

- A. Waist circumference is the most common tool, as it directly relates to central body fat risks.
 - B. Body Mass Index (BMI) is the most commonly used tool, but there's a shift due to its inaccuracy for certain populations.
 - C. Body fat percentage is the most common due to its detailed insights, but BMI is gaining popularity.
 - D. The Edmonton obesity staging system is most commonly used, but waist circumference is being considered more often.
-

2. In the context of obesity, which of the following genetic factors would most likely contribute to increased hunger sensations?

- A. An imbalance in the production of insulin.
 - B. Variations causing leptin resistance.
 - C. Decreased secretion of cholecystokinin.
 - D. Increased production of pancreatic polypeptide.
-

3. Which of the following scenarios best illustrates an environmental risk factor contributing to obesity?

- A. An individual with genetic variations affecting hormone secretion.
 - B. Someone who experiences long-term financial stress without access to nutritious food.
 - C. An individual with a sedentary lifestyle due to lack of motivation.
 - D. A person with high levels of ghrelin due to a hormonal condition.
-

4. How might the gut microbiome indirectly contribute to obesity, based on course material?

- A. By causing direct increases in ghrelin secretion leading to enhanced hunger.
 - B. By affecting the body's thermal energy balance and increasing energy expenditure.
 - C. By influencing systemic inflammation through diet, affecting insulin sensitivity.
 - D. By directly reducing leptin levels, leading to loss of appetite regulation.
-

5. Which of the following statements accurately describes the limitations of BMI in diagnosing obesity as mentioned in the course content?

- A. BMI does not account for body fat distribution, leading to inaccurate cardiac risk assessments.

- B. BMI is a sufficient tool for Asians but is less reliable for older adults due to muscle mass loss.
 - C. BMI is not applicable for pediatric populations who require waist circumference evaluations.
 - D. BMI provides reliable results across all populations, making alternative measures unnecessary.
-

6. Which of the following best describes the interrelationship between the biomechanical and metabolic conditions of obesity?

- A. Biomechanical conditions are caused by increased metabolism due to obesity.
 - B. Metabolic conditions exclusively lead to biomechanical disorders.
 - C. Biomechanical conditions can influence metabolism, and metabolic conditions can affect biomechanics.
 - D. Metabolic conditions occur as a result of biomechanical changes, not the other way around.
-

7. What is the primary mechanism by which obesity contributes to type 2 diabetes?

- A. It directly increases glucose production in the liver.
 - B. Obesity causes insulin resistance mainly due to inflammation from non-esterified fatty acids.
 - C. Obesity decreases insulin secretion by damaging pancreatic cells.
 - D. Type 2 diabetes only occurs in the presence of both obesity and hypertension.
-

8. When it comes to pharmacological treatment options for obesity, how do GLP-1 receptor agonists function?

- A. They work by increasing lipolysis and reducing fat storage in cells.
 - B. They inhibit fat absorption by the digestive tract.
 - C. They slow gastric emptying and affect the central nervous system to control appetite.
 - D. They enhance the metabolism of glucose by increasing pancreatic enzyme activity.
-

9. Which statement correctly differentiates the risks of coronary artery disease in individuals with obesity?

- A. Coronary artery disease is directly caused by obesity in all cases.
 - B. As BMI increases, the risk for coronary artery disease increases due to the presence of multiple factors.
 - C. Obesity is unrelated to coronary artery disease, which is primarily due to lifestyle choices.
 - D. The risk is determined mostly by the genetic predisposition of the individual rather than obesity itself.
-

10. Why might lifestyle changes be considered the first line of treatment in obesity management over surgical options?

- A. Lifestyle changes have been shown to be dramatically more effective than surgeries in short-term weight loss.

- B. Surgery is more risky and expensive, making lifestyle changes safer and more accessible for many individuals.
 - C. Surgical procedures do not address the underlying causes of obesity like lifestyle changes do.
 - D. There are fewer post-operative complications with lifestyle changes compared to surgeries.
-

Copyright © 2025 Cheap Nursing CEUs

Visit us at <https://www.cheapnursingceus.com>