

# Cheap Nursing CEUs

## Pediatric Acute Respiratory Infection

**1. Which of the following best represents a risk factor for acute respiratory infections for pediatric patients?**

- A. Weight loss
- B. A body mass index of 24.9 kg/m<sup>2</sup>
- C. Mobility limitations
- D. Passive smoking

---

**2. Which of the following best represents a sign/symptom of an acute respiratory infection?**

- A. Nasal congestion
- B. Oral dysphagia
- C. Polyphagia
- D. Polydipsia

---

**3. A health care professional has questions regarding respiratory infections. Which of the following informational points of interest should be communicated to the health care professional?**

- A. Respiratory infections are typically only caused by viruses.
- B. Respiratory infections are typically only caused by bacteria.
- C. The leading causes of acute respiratory infections include viruses and bacteria.
- D. The leading causes of acute respiratory infections include viruses and extreme weight loss.

---

**4. The parents of a 2-year-old child have questions regarding the common cold. Which of the following educational points should be expressed to the parents?**

- A. Most individuals recover from the common cold within, approximately, 7 - 10 days.
- B. Most individuals recover from the common cold within, approximately, 14 - 21 days.
- C. Treatment for the common cold typically involves antibiotics.
- D. Treatment for the common cold typically involves antibiotics and acetaminophen.

---

**5. Which of the following medications is recommended as first-line therapy for acute bacterial sinusitis?**

- A. Amoxicillin
- B. Cephalexin
- C. Cefadroxil
- D. Clindamycin

---

**6. A health care professional is developing an educational lecture centered around bronchiolitis. Which of the following informational points of interest should be included in the health care professional's lecture?**

- A. Bronchiolitis is typically caused by the RSV.
- B. Transmission of RSV typically only occurs through contaminated food.
- C. Individuals infected with RSV are usually contagious for three to eight weeks.
- D. Early symptoms of RSV may include weight gain.

---

**7. Which of the following statements is most accurate?**

- A. In very young infants (e.g., older than six months old), the only symptoms of a RSV infection that may be present include the following: irritability, weight loss, and odynophagia.
- B. In very young infants (e.g., older than six months old), the only symptoms of a RSV infection that may be present include the following: irritability, increased appetite, and apnea.
- C. In very young infants (e.g., less than six months old), the only symptoms of a RSV infection that may be present include the following: irritability, otalgia, and odynophagia.
- D. In very young infants (e.g., less than six months old), the only symptoms of a RSV infection that may be present include the following: irritability, decreased appetite, and apnea.

---

**8. Infants/children suffering from bronchitis should seek health care if they meet which of the following conditions?**

- A. Symptoms lasting more than one week
- B. Symptoms lasting more than two weeks
- C. The child is younger than two months old and has a fever of 102.4 °F
- D. The child is younger than three months old and has a fever of 100.4 °F

---

**9. A 10-year-old patient presents to a health care facility. The patient is diagnosed with community-acquired pneumonia. The patient's parents report the patient has a hypersensitivity to penicillin. Which of the following medications should be used to treat the aforementioned patient?**

- A. Amoxicillin
- B. Azithromycin
- C. Peramivir
- D. Remdesivir

---

**10. A health care professional has questions regarding the differences between influenza and a "cold." Which of the following informational points of interest should be communicated to the health care professional?**

- A. Symptom onset is abrupt with a cold.
- B. Symptom onset is abrupt with influenza.
- C. Chills are common with a cold and not as common with influenza.
- D. Headaches are common with a cold and less common with influenza.

---

**11. Which of the following medications is approved for treatment of influenza in children two weeks old or older?**

- A. Oseltamivir
- B. Zanamivir
- C. Peramivir
- D. Baloxavir

---

**12. Which of the following best represents a COVID-19 treatment recommendation for children?**

- A. Most children with SARS-CoV-2 infection will require amoxicillin.
- B. Most children with SARS-CoV-2 infection will require cephalexin.
- C. Remdesivir is recommended for hospitalized children aged  $\geq$ 12 years with COVID-19 who have risk factors for severe disease and have an emergent or increasing need for supplemental oxygen.
- D. Remdesivir is recommended for hospitalized children aged  $\geq$ 14 years with COVID-19 who have risk factors for severe disease and do not have an emergent or increasing need for supplemental oxygen.

---

**13. A health care professional has questions regarding mastoiditis. Which of the following informational points of interest should be communicated to the health care professional?**

- A. Mastoiditis is often associated with the common cold.
- B. Mastoiditis is often associated with influenza.
- C. Symptoms of mastoiditis include nasal congestion and oral dysphagia.
- D. Symptoms of mastoiditis include: fever, pain, ear redness, and discharge from the ear.

---

**14. Which of the following best represents a sign/symptom of MIS-C?**

- A. Extremely high blood pressure
- B. Low blood pressure
- C. Weight gain
- D. Weight gain with increased appetite

---

**15. The parents of a newborn child have questions regarding adequate nutrition. Which of the following educational points should be expressed to the parents?**

- A. Individuals should only exclusively feed infants human milk for the first two months of life.
- B. Individuals should exclusively feed infants human milk for the first six months of life.
- C. Individuals should provide infants with supplemental vitamin D beginning four months after birth.
- D. Individuals should provide infants with supplemental vitamin D beginning six months after birth.

---

**16. Which of the following statements is most accurate?**

- A. Inactivated vaccines include the dead version of the infectious agent that causes a specific infection or disease.
- B. Inactivated vaccines include the live version of the infectious agent that causes a specific infection or disease.
- C. Live-attenuated vaccines are not similar to the natural infection they help prevent - therefore, they create a long-lasting immune response of about six months.
- D. Live-attenuated vaccines are similar to the natural infection they help prevent - therefore, they create a short-term immune response.

---

**17. A health care professional has questions regarding Afluria Quadrivalent. Which of the following informational points of interest should be communicated to the health care professional?**

- A. Afluria Quadrivalent is approved for use in individuals six months of age and older.
- B. Afluria Quadrivalent is only approved for use in individuals six years of age and older.
- C. The recommended dose for Afluria Quadrivalent is 0.25 mL.
- D. The recommended dose for Afluria Quadrivalent is 1.5 mL.

---

**18. What is the recommended dose for FluMist?**

- A. 0.2 mL
- B. 0.25 mL
- C. 0.5 mL
- D. 1.5 mL

---

**19. Which of the following statements regarding Prevnar 13 is most accurate?**

- A. Children six weeks through five years should receive a three-dose immunization series consisting of a 0.5 mL intramuscular injection administered at 4, 6, and 12 - 15 months of age.
- B. Children six weeks through five years should receive a four-dose immunization series consisting of a 0.5 mL intramuscular injection administered at 2, 4, 6, and 12 - 15 months of age.
- C. Children six through 17 years of age and adults should receive a 2-dose immunization series over the course of four months.
- D. Children six through 17 years of age and adults should receive a 2-dose immunization series over the course of eight months.

---

**20. A health care professional has questions regarding Pneumovax 23. Which of the following informational points of interest should be communicated to the health care professional?**

- A. Pneumovax 23 should be administered in a single oral dose.
- B. Pneumovax 23 should be administered in a single dose, either by intramuscular injection or subcutaneous injection.
- C. Pneumovax 23 should be administered in a 2-dose immunization series.

D. Pneumovax 23 should be administered in a 4-dose immunization series.

---

Copyright © 2026 Cheap Nursing CEUs

Visit us at <https://www.cheapnursingceus.com>