

# Cheap Nursing CEUs

## Sexually Transmitted Infections (STIs)

**1. Which of the following factors is primarily associated with the increased risk of STI transmission among adolescents?**

- A. Frequent international travel
  - B. High socioeconomic status
  - C. Engagement in survival sex
  - D. Limited access to cultural activities
- 

**2. Chlamydia is often asymptomatic, but when symptoms do occur, which of the following is NOT typically associated with this STI?**

- A. Dysuria
  - B. Pain during bowel movements
  - C. Vaginal discharge
  - D. Testicular pain
- 

**3. Regarding the screening guidelines for Hepatitis B, which of the following populations should be prioritized?**

- A. All sexually active men over 40
  - B. Men with a history of syphilis
  - C. Pregnant women
  - D. Adolescents with no symptoms
- 

**4. Why is it crucial for healthcare providers to distinguish between behaviors and biological risk factors when assessing STI risk?**

- A. To ensure patients have access to social services
  - B. To implement appropriate preventive measures and screenings
  - C. To prioritize treatment for respiratory infections
  - D. To avoid unnecessary STI testing
- 

**5. Which statement is true about the implications of untreated syphilis?**

- A. It has no impact on the nervous system
  - B. It results exclusively in ocular damage
  - C. Untreated syphilis can lead to tertiary complications including organ damage
  - D. It can cause bone density issues exclusively
-

**6. According to the 2021 updated guidelines, why is rectal chlamydia testing recommended over azithromycin for treating rectal infections?**

- A. Azithromycin is less effective than doxycycline for rectal infections.
  - B. Doxycycline causes fewer side effects than azithromycin for rectal infections.
  - C. Azithromycin has more resistance issues with rectal infections.
  - D. Doxycycline is preferred due to quicker symptom relief.
- 

**7. What is the primary reason for recommending hepatitis B vaccination for MSM who are not already vaccinated?**

- A. Higher prevalence of hepatitis B in MSM compared to other groups.
  - B. Increased risk of exposure due to higher rates of unprotected sex.
  - C. Lack of routine testing makes vaccination crucial.
  - D. Enhanced efficacy of vaccination in preventing hepatitis B complications.
- 

**8. How can nurses create a non-judgmental clinical environment when discussing sexual health with patients?**

- A. By ensuring questions about sexual practices are asked to certain patients based on visible risk factors.
  - B. By asking questions to all patients solely based on age and sexual orientation.
  - C. By explaining that the same questions are asked to every patient, regardless of background.
  - D. By avoiding direct questions about the number of sexual partners.
- 

**9. Which of the following best describes the implications of untreated STIs as highlighted in the course?**

- A. Untreated STIs can lead only to local symptoms without systemic impact.
  - B. Untreated STIs primarily cause psychological issues and rarely physical ones.
  - C. Untreated STIs can cause serious health issues, including infertility and neurological diseases.
  - D. Untreated STIs are mostly an inconvenience with few long-term effects.
- 

**10. In Section 7, what is emphasized as a crucial component of STI prevention through nursing education?**

- A. Promoting abstinence as the only sure method of prevention.
  - B. Encouraging monogamous relationships as a way to prevent STIs.
  - C. Providing accurate disease information and encouraging regular testing.
  - D. Recommending vaccination as the primary prevention strategy.
-

Visit us at <https://www.cheapnursingceus.com>